

Toward a synthetic understanding of the role of phenology in ecology and evolution

Jessica Forrest and Abraham J. Miller-Rushing

Phil. Trans. R. Soc. B 2010 **365**, 3101-3112
doi: 10.1098/rstb.2010.0145

References

This article cites 160 articles, 33 of which can be accessed free
<http://rstb.royalsocietypublishing.org/content/365/1555/3101.full.html#ref-list-1>

Rapid response

[Respond to this article](#)
<http://rstb.royalsocietypublishing.org/letters/submit/royptb;365/1555/3101>

Subject collections

Articles on similar topics can be found in the following collections

[ecology](#) (2301 articles)
[evolution](#) (2591 articles)

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

To subscribe to *Phil. Trans. R. Soc. B* go to: <http://rstb.royalsocietypublishing.org/subscriptions>

Introduction

Toward a synthetic understanding of the role of phenology in ecology and evolution

Jessica Forrest^{1,2,*} and Abraham J. Miller-Rushing^{3,4,5}

¹*Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada M5S 3G5*

²*Rocky Mountain Biological Laboratory, Crested Butte, CO 81224, USA*

³*USA National Phenology Network, Tucson, AZ 85719-5224, USA*

⁴*The Wildlife Society, Bethesda, MD 20814-2197, USA*

⁵*National Park Service, Acadia National Park, Schoodic Education and Research Center, Bar Harbor, ME 04609, USA*

Phenology affects nearly all aspects of ecology and evolution. Virtually all biological phenomena—from individual physiology to interspecific relationships to global nutrient fluxes—have annual cycles and are influenced by the timing of abiotic events. Recent years have seen a surge of interest in this topic, as an increasing number of studies document phenological responses to climate change. Much recent research has addressed the genetic controls on phenology, modelling techniques and ecosystem-level and evolutionary consequences of phenological change. To date, however, these efforts have tended to proceed independently. Here, we bring together some of these disparate lines of inquiry to clarify vocabulary, facilitate comparisons among habitat types and promote the integration of ideas and methodologies across different disciplines and scales. We discuss the relationship between phenology and life history, the distinction between organismal- and population-level perspectives on phenology and the influence of phenology on evolutionary processes, communities and ecosystems. Future work should focus on linking ecological and physiological aspects of phenology, understanding the demographic effects of phenological change and explicitly accounting for seasonality and phenology in forecasts of ecological and evolutionary responses to climate change.

Keywords: climate change; life history; natural selection; phenology; synchrony

1. INTRODUCTION

The word ‘phenology’ has the same Greek root, *phainomai* (‘to appear’), as the words ‘phenomenon’ and ‘phenotype’. Although the latter two words may be more familiar, phenology—the study of the timing of recurring seasonal biological events—has existed as a field of scientific inquiry for centuries. Whether for agricultural or religious reasons, or simply as a way of marking the passage of the seasons, humans have long had an interest in documenting the more-or-less regular appearances of such things as the first flower blossoms of spring, the first migrating birds or the first frost-damaged leaves of fall (Hopkins 1918; Sparks & Menzel 2002; Aono & Kazui 2008). The ancient Greeks themselves recognized the value of phenology—a more reliable indicator of local weather than the movement of the constellations—and used the timing of leaf fall as a guide for when to sow winter crops (Bostock & Riley 1855).

At its simplest, phenology is merely the temporal dimension of natural history. However, this temporal dimension is critical, because it determines the stage of development reached by an organism or population

at the time when it intersects with particular components of its environment. Phenology is therefore a major structuring element in nearly all areas of ecology and evolution. Historically, because of its practical importance for plant cultivation, much phenological research has focused on agricultural applications such as pest management, agricultural meteorology and horticulture (Hopkins 1918; Garner & Allard 1920; Schwartz *et al.* 1997). Ecological and evolutionary studies with a focus on phenology also have a long history (e.g. Robertson 1924; Leopold & Jones 1947); however, many studies with important phenological components did not refer to these as phenology *per se* (e.g. Clausen *et al.* 1941; Corbet 1954; Janzen 1967).

In the last two decades, growing concern with documenting and forecasting the impacts of climate change has driven increased interest in the role of phenology in ecology and evolution. Phenological shifts have been among the most obvious and thoroughly documented biological responses to the climate warming of the last 150 years (Beebe 1995; Myneni *et al.* 1997; Crick & Sparks 1999; Fitter & Fitter 2002; Parmesan & Yohe 2003). At the same time, progress in elucidating the genetic basis of flowering time in plants (Ausin *et al.* 2005; Buckler *et al.* 2009; Wang *et al.* 2009), diapause induction in insects (Tauber *et al.* 2007) and offspring hatching date in birds (Liedvogel *et al.* 2009) is bringing a more

* Author for correspondence (jessica.forrest@utoronto.ca).

One contribution of 11 to a Theme Issue ‘The role of phenology in ecology and evolution’.

mechanistic understanding of phenology within reach. Advances in the fields of molecular and developmental biology, quantitative genetics, phylogenetics and ecosystem ecology have also contributed to the recent growth of phenological research.

To date, these various strands of phenology research have tended to proceed independently and have employed different terminologies. In convening this themed issue, we aim to bring together some of these disparate lines of inquiry to clarify vocabulary, facilitate comparisons among habitat types and, most of all, promote the integration of ideas and methodologies across different disciplines and scales. This issue also emphasizes the importance of phenology in nearly all aspects of ecology and evolution.

In this introduction, we start by clarifying the relationship between phenology and life history, and by briefly reviewing the physiological processes and environmental cues governing phenology in different taxa. We then move from the individual organism to the level of the population, and discuss how the shape of the population-level phenological distribution can be characterized—and why it matters. Finally, we provide an overview of the role of phenology in the ecology of communities and ecosystems, and in the evolution of adaptation (or, sometimes, maladaptation). Along the way, we outline some of the main challenges and areas for further work in this field.

2. PHENOLOGY AND LIFE HISTORY

Historical observations of phenology, as well as many recent ecological studies, have mainly concerned patterns at the population level. These studies ask questions such as: how does a plant population's timing of leaf unfolding vary with respect to temperature? Or, has the date of the first frog call advanced over a period of decades? At the individual level, in contrast, the question of interest might be: why does an individual of a particular size or sex begin growth or reproduction at a given time of year? Individual-level patterns are less often equated with phenology (Visser *et al.* 2010), but understanding them is essential for making sense of many population-level patterns, which, after all, represent the integrated activity schedules of many individuals.

The term phenology is sometimes used interchangeably with life history because both incorporate the timing of growth, reproduction and senescence. Of course, phenology does not encompass such non-temporal aspects of life history as size at reproductive maturity and brood size. However, interpreting phenology in the context of life history allows us to integrate phenological investigations with the existing theory and experiments that describe life-history evolution—e.g. the trade-offs that underlie why, in an ultimate sense, annual plants flower at a particular time or why tadpoles metamorphose when they do. Unfortunately, life-history theory and the implications of relevant trade-offs are rarely included in studies exploring variation among species in recent shifts in phenology (Fitter *et al.* 1995; Bradley *et al.* 1999; Miller-Rushing & Primack 2008).

One such trade-off occurs between optimal age (young) and size (large) at maturity. The realized

life-history strategy of an individual is expected to reflect some balance between these, with the exact point of compromise influenced by factors such as sex of the individual (Morbey & Ydenberg 2001; Nève & Singer 2008) or relative risk of mortality in larval and adult habitats (Werner 1986; Abrams & Rowe 1996). Environmental factors can obscure the trade-off: individuals growing in a high-quality environment can both be large at maturity and reach maturity early (van Noordwijk & de Jong 1986). Nevertheless, the age–size compromise may influence the type or magnitude of evolutionary change in phenology that would be expected in response to a warming climate (Etterson & Shaw 2001). In annual plants, for example, there is frequently a positive genetic correlation between age and size at flowering (Mitchell-Olds 1996; Franks & Weis 2008); in insects, many of which are likewise annuals, later metamorphosis to adulthood means more time for growth (Masaki 1967). In both cases, the optimal phenological response to an extended growing season depends on the relative benefits of reaching reproductive maturity earlier in the season or growing larger before reproducing.

The utility of this basic life-history framework depends on how a species' lifespan and schedule of reproduction fit within the annual cycle. The expected trade-off between optimal timing and size at reproduction is modified in iteroparous species, which can use resources acquired in a previous growing season for reproduction in the current year. For this reason, in temperate-zone perennials, large plants frequently flower earlier than smaller individuals in the same populations (Forrest & Thomson 2010 and references therein). Similarly, birds in good condition generally lay eggs earlier in a given season than those in poor condition (Price *et al.* 1988; Rowe *et al.* 1994). In red squirrels (*Tamiasciurus hudsonicus*), variation in breeding date is also influenced by maternal condition, itself a response to food availability in the previous year (Réale *et al.* 2003). Thus, both iteroparity and environmental variation in resource availability obscure the time–size trade-off because the resources available for reproduction are not solely determined by the individual's ability to acquire resources in a particular growing season.

In addition, many species do not reproduce on an annual schedule. The phenology of flowering and fruiting in southeast Asian rain forests, in which community-wide mass-flowering events take place at irregular intervals of more than 1 year (Medway 1972; Brearley *et al.* 2007), has little to do with the life history of individual trees and much to do with the factors favouring population- and community-level synchrony among individuals. In short-lived taxa with several generations per year, the link between life history and phenology may likewise not be immediately apparent. However, life-history theory can still be useful; for example, the expected number of generations per year in multi-voltine insects—and, therefore, the times of year when particular life stages will be abundant—is the outcome of the same age–size optimization problem described above (Roff 1980). In general, integrating life-history theory into ecological studies should help both in forecasting

changes in phenology and in understanding whether the changes observed so far are likely to be adaptive responses to new conditions.

3. THE MECHANISTIC BASIS OF PHENOLOGY

Just as it is important to understand the life-history trade-offs that are the ultimate causes of many phenological patterns, understanding the proximate drivers of phenology is critical if we wish to predict phenological responses to environmental change. Forecasts of evolutionary change in phenology based on simple optimality models could well be modified by an understanding of the genetics and physiology involved, including the pleiotropic effects of alleles affecting phenological traits (Metcalf & Mitchell-Olds 2009). The timing of many phenological events (e.g. onset of reproduction, entry into or emergence from a dormant stage) results proximally from a complex interplay among an organism's genes and several external environmental factors. These environmental factors, such as temperature or precipitation (see below), may directly control the timing of biological events, or they may act instead as cues that set the organism's internal 'biological clock' (Gwinner 1996; Ausin *et al.* 2005). For most species, however, we do not know (i) the specific environmental factors that are most important in determining phenology, (ii) the precise molecular and physiological processes that regulate phenology, and (iii) whether variation in phenology over time or among individuals reflects genetic differences or simply plastic responses to environmental heterogeneity. Rapid progress is being made to address these uncertainties about the mechanisms regulating phenology (e.g. Visser *et al.* 2010; Wilczek *et al.* 2010), but for now, they substantially limit our ability to anticipate future responses to changes in a variety of climate variables. Here, we review some of the best-studied factors that are known to affect the phenology of plants and animals.

(a) Genes

Some of the variation in phenological traits between individuals and populations clearly has a genetic basis. This conclusion is supported by heritability estimates (reviewed by Mazer & LeBuhn 1999; Geber & Griffen 2003; Hendry & Day 2005) as well as empirical demonstrations of evolution in phenological traits (Paterniani 1969; Réale *et al.* 2003; Bradshaw & Holzapfel 2006; Franks *et al.* 2007). Genes may confer a propensity for earlier growth or reproduction regardless of environmental conditions, or they may affect an individual's sensitivity to the environmental conditions that affect timing. For example, in *Arabidopsis thaliana*, plants carrying different alleles at the *FRIGIDA* or *PHYC* loci differ in their sensitivity to vernalization or photoperiod, respectively, and therefore in the relationship between flowering time and environment (Stinchcombe *et al.* 2004; Balasubramanian *et al.* 2006). Similarly, variants of the *timeless* gene differentially affect sensitivity to diapause cues in certain European populations of *Drosophila melanogaster* (Tauber *et al.* 2007).

(b) Photoperiod

In several cases where the genetic basis of phenological traits has been confirmed, the alleles involved confer different levels of responsiveness to photoperiod cues (Bradshaw & Holzapfel 2001; Sandrelli *et al.* 2007; Van Dijk & Hautekèete 2007). The predictability of the seasonal change in the light–dark cycle at a given latitude makes photoperiod a reliable indicator of the time of year, at least away from the equator; accordingly, many organisms use changing daylength as a cue for the initiation of reproduction, hibernation, migration, diapause or moult. Photoperiod plays a role in regulating seasonal patterns in such distantly related organisms as mustards (Ausin *et al.* 2005), mollusks (Wayne 2001) and mammals (Goldman 2001). Changing daylength influences the timing of sexual reproduction in some freshwater zooplankton (Stross & Hill 1968; Gilbert 1974), the timing of spore germination in marine diatoms (Eilertsen *et al.* 1995) and the induction and termination of diapause in freshwater copepods (Williams-Howze 1997). Seasonal variation in insolation may even influence the timing of leaf flush in 'aseasonal' tropical forests (van Schaik *et al.* 1993). In many insects, diapause initiation and—perhaps less commonly—termination are controlled by photoperiod, although the resumption of activity following diapause is likely to depend on other factors, such as temperature, as well (Mazaki 1980; Tauber *et al.* 1986).

(c) Temperature

Dependence on photoperiod cues alone would render organisms vulnerable to mistiming their activities in years with unusual weather conditions, or in the event of rapid climate change. However, the photoperiod response is often modified or even overridden by other, more directly relevant environmental factors, of which the most commonly used, at least in temperate climates, is temperature. The interaction between long days and warm temperatures has been well characterized in the flowering pathway of *A. thaliana* (Ausin *et al.* 2005; Wilczek *et al.* 2009). Certain migrating birds also integrate information on temperature and photoperiod (Bauer *et al.* 2008).

Although endotherms such as birds may use temperature, like photoperiod, simply as a cue informing them of the likely future availability of food, in other organisms, temperature affects phenology directly by influencing the rates of biochemical processes (cf. Gillooly *et al.* 2002). As a consequence, the accumulation of a certain number of heating units (e.g. degree-days) often predicts well the date of flowering in plants (e.g. Jackson 1966; Diekmann 1996), and flowering phenology commonly tracks interannual variation in air temperatures (Fitter *et al.* 1995; Sparks *et al.* 2000; Miller-Rushing *et al.* 2007). Heat accumulation similarly affects development rate and, hence, the timing of appearance of adults, in many economically important insect species (Embree 1970; Kemp & Onsager 1986; Kemp *et al.* 1986; Régnière *et al.* 2007). In multi-voltine insects (those with multiple generations in a year), shortening days late in the year commonly induce diapause regardless of

temperature; but the number of generations achieved prior to this will depend on temperatures experienced, and hence the rate of development, up to that point (Tobin *et al.* 2008).

However, there is often more to the temperature effect than simple heat accumulation. Many plants have a chilling requirement, such that subsequent development is delayed or prevented if they have not experienced cold winter temperatures (Murray *et al.* 1989; Morin *et al.* 2009). This requirement is referred to as vernalization when applied to flowering (Henderson *et al.* 2003). The need for cool temperatures has the counterintuitive effect of delaying phenology in warm years (Zhang *et al.* 2007). A similar phenomenon has been documented in insects: in several temperate-zone species, a longer overwintering period reduces the heat requirement for springtime emergence of adults (Kimberling & Miller 1988; Bosch & Kemp 2003, 2004). Other factors that complicate the relationship between temperature and phenology are differences between species in their lower threshold temperatures for development (Kemp & Dennis 1989) or in their abilities to behaviourally thermoregulate by moving into patches of sunlight or shade (van Nouhuys & Lei 2004).

(d) *Precipitation*

In the tropics and arid environments, variation in precipitation is more likely than temperature to drive phenological patterns. In different types of tropical forests, either rain or drought can induce flowering (Medway 1972; van Schaik *et al.* 1993; Brearley *et al.* 2007); often, this does not occur on an annual cycle. Many desert plants germinate (annuals) or resume growth (perennials) in response to rainfall (Beatley 1974; Zhang *et al.* 2006; Kimball *et al.* 2010). Desert animals often emerge from diapause or aestivation in response to moisture (Cloudsley-Thompson 1991; Danforth 1999).

At high altitudes and latitudes, flowering time and insect activity can be strongly, and apparently linearly, correlated with timing of snowmelt (Ellebjerg *et al.* 2008; Høye & Forchhammer 2008; Forrest *et al.* 2010). However, it is not clear that snowmelt is itself a cue to which organisms respond. Instead, disappearance of snowpack may set a lower bound on the date at which heat units can begin to accumulate (Thórhallsdóttir 1998). Thus, extremely early snowmelt unaccompanied by warm early-spring temperatures—a conjunction of circumstances that can occur if there is little snowfall the previous winter—may fail to advance phenology. This can look like an accelerating relationship between phenology and snowmelt date (Inouye 2008; Steltzer *et al.* 2009), but a simple, uniform response to accumulated degree-days in a given year may be a more parsimonious interpretation.

Often it is not possible to compare alternative environmental predictors of phenology (e.g. temperature versus snowmelt) because detailed weather records are not available. This illustrates a common limitation of descriptive phenological studies: it is relatively easy to detect a correlation between some climate variable and a particular phenological

response; but this in itself does not demonstrate that the climate variable in question is the proximate cue regulating phenology. This is simply another case of correlation not equalling causation: multiple climate factors are likely to covary, and standard experimental designs (such as snow removal or warming structures), while valuable in their own right, may be inadequate for separating these variables. More tightly controlled experiments are necessary to determine unequivocally which environmental factors regulate phenology (e.g. Cleland *et al.* 2006; Sherry *et al.* 2007). Where experiments are impossible, statistical modelling to compare the effectiveness of different predictors can at least provide clues about which cues are most likely involved (e.g. Dunne *et al.* 2003; Hülber *et al.* 2010). Similar responses to recent climate change among groups of related species (i.e. phylogenetic conservatism in phenological shifts; Davis *et al.* 2010) suggest common drivers of phenology within clades; this may permit inferences about mechanism in taxa that have not yet been studied. A better mechanistic understanding is necessary if we are to make predictions about phenological responses to future, novel climates, and the chances of phenological decoupling among interacting species (see below; Araújo & Luoto 2007).

4. CHARACTERIZING PHENOLOGIES AT THE POPULATION LEVEL

The ultimate and proximate factors that regulate the phenologies of individual organisms contribute in turn to phenological patterns at the level of the population or community. Phenology, as a characteristic of the population, has the components of any statistical distribution. In the case of flowering phenology, these include the mean flowering date, duration (range) of flowering and the higher moments such as variance and skewness. Importantly, population-level distributions can be inferred from the traits of individuals, but not always vice versa: a skewed flowering distribution could result from individuals having skewed flowering curves, or from individuals with symmetrical flowering curves having a skewed distribution of first flowering dates. The positions of phenological distributions (i.e. first dates, means or peaks) have received the most attention from population and community ecologists, because of the consequences for overlap with other temporally varying components of the environment (see below; Araújo & Luoto 2007). However, the distributions as a whole, and precisely how they relate to individual-level phenology, have received less attention to date (though see Laaksonen *et al.* 2006; Elzinga *et al.* 2007). This lack of attention—caused in part by the rarity of adequate datasets—limits our ability to understand the ecological and evolutionary consequences of population-level phenology, including the availability of temporal niches for non-native species (Wolkovich & Cleland *in press*) and the form of selection on phenological traits. Here, we outline some aspects of phenological distributions that are often overlooked.

Variance and kurtosis ('peakedness') in phenology reflect within-population synchrony. Synchrony in reproduction can improve chances of mate-finding

(Augspurger 1981; Reed *et al.* 2009) and offspring survival (Ims 1990; Kelly & Sork 2002), but it also increases competition for resources. Less obviously, decreased variance in phenology at one trophic level can affect higher trophic levels, which may depend on the food supply being more evenly distributed in time—that is, having higher among-individual or among-plot variability in phenological events (Post *et al.* 2008). Changes in population and community-level synchrony in response to warming temperatures are worth monitoring because of these potential effects on demography and ecosystem processes (Miller-Rushing *et al.* 2010).

Skewness is common in phenological distributions, and it determines the extent to which the population mean is an adequate reflection of central tendency: in strongly skewed distributions, shifts in the mean, rather than the median, poorly represent trends experienced by most individuals. Timing of germination and flowering in plant populations is often positively skewed (Rabinowitz *et al.* 1981; Rathcke & Lacey 1985; Brown & Mayer 1988), as are timing of emergence in insects (Danks 2006) and arrival and laying dates in migratory birds (Sparks *et al.* 2005; Laaksonen *et al.* 2006). This pattern may arise because most individuals respond rapidly and similarly to the relevant environmental cues, while a smaller number experience problems in development or migration that delay phenology to varying extents (Rathcke & Lacey 1985; Danks 2006). Intriguingly, skewness often increases in warm years, with populations developing a longer tail at the end of the season (Roy & Sparks 2000; Sparks *et al.* 2005; Forrest & Thomson 2010). Skewness also determines the extent to which an individual's timing of activity covaries with population density. This makes directional selection on temporal traits difficult to distinguish from stabilizing selection: given a positively skewed distribution, selection for earliness resembles selection for synchrony.

Although recognition of the full shape of phenological distributions is important for many ecological and evolutionary questions, a framework based on a simple, unimodal trait distribution will be inadequate for characterizing some cyclical phenomena. Primary production in many aquatic habitats, for example, does not have a clearly defined duration or even, in some cases, an obvious seasonal peak. Simply demonstrating the frequency and consistency of population or community cycles—a prerequisite for documenting effects of climate change on phenology—can be a challenge in such systems. Winder & Cloern (2010) overcome this challenge with an innovative approach: wavelet analysis applied to time series of phytoplankton biomass. Elsewhere, Altermatt (2010) has used a kernel-density estimation function to describe the multi-modal distributions produced by multi-voltine insects. Techniques like these could have broad applicability to systems where analysing temporal trends in phenology would otherwise be problematic.

5. ECOLOGICAL EFFECTS OF PHENOLOGY

There has been much attention in recent literature to the likely ecological consequences of shifts in

phenological distributions in response to climate change. Because phenology is involved in nearly all ecological relationships, there is clearly potential for important effects. Here, we highlight a few of the consequences of changing phenology for population dynamics, species interactions and ecosystems.

One often-discussed possible result of climate change is that species will differ in the degree to which their phenologies shift, with potentially dire consequences for interacting species (e.g. Harrington *et al.* 1999; Durant *et al.* 2007; Both *et al.* 2009; Hegland *et al.* 2009). In principle, these shifts could have positive or negative consequences for the populations involved, depending on whether the interaction in question is mutually beneficial (i.e. a mutualism), mutually detrimental (i.e. competition) or unilaterally beneficial (e.g. predation), and whether differential changes in phenology drive species closer together or further apart in time. Such shifts in interactions seem inevitable, insofar as species use different cues to regulate phenology (still something of an unanswered question; Aono & Kazui 2008; see above). In practice, however, there are still few examples of such shifts having detectable demographic consequences. Reasons for this persistent gap, and possible solutions, are discussed by Miller-Rushing *et al.* (2010). A convincing demonstration requires showing that a change in interaction strength or frequency has occurred, that this change is the result of climate change and that the change has altered the vital rates of one or more of the species involved. In this issue, Thomson (2010) provides one of the few examples of an important species interaction that has been documented over the long term, showing that pollen limitation in a subalpine wildflower has increased over the last 17 years, and suggesting that plant–pollinator decoupling may be occurring. This is a phenomenon that has been predicted by many (e.g. Dunne *et al.* 2003; Memmott *et al.* 2007), but not previously demonstrated. However, the data are inconclusive as to whether climate change is responsible, and population declines have yet to be observed: the plant is a perennial, and we do not know whether population size is limited by seed supply. This illustrates the difficulties inherent in this type of work and suggests where further efforts are required.

Shifts in the timing of reproduction, in particular, have possible consequences beyond changing species interactions. The need to fit at least one reproductive episode into the annual cycle can be the factor limiting a species' geographical range (e.g. Jönsson *et al.* 2009), such that longer growing seasons can allow species establishment beyond the current range limit. This is the rationale behind process-based models such as PHENOFIT, described in this issue by Chuine (2010). Such approaches promise more mechanistically grounded forecasts of species range changes with climate warming than have been provided by purely correlation-based 'climate envelope' techniques. Furthermore, for some species, early completion of a first bout of reproduction may permit a second breeding attempt in the same season. This is particularly likely if the tail end of the growing season is being extended as well. Several

temperate-zone birds and other taxa are capable of double-brooding if there is time (Verhulst *et al.* 1997 and references therein; Saino *et al.* 2004), provided food resources are also sufficient (cf. Husby *et al.* 2009). In short-lived species, warmer temperatures and longer growing seasons may allow additional generations per year (e.g. Tobin *et al.* 2008; Jönsson *et al.* 2009; Altermatt 2010). Both of these possibilities have major implications for population growth. However, because not all species are capable of multiple broods or generations in a year, even when growing season length is adequate, some species will benefit more than others from warming. In particular, there is concern that outbreaks of certain insect pests will increase in frequency (Logan *et al.* 2003). Interestingly, this is not only an ecological advantage; in principle, having more generations in a given time span could allow more rapid adaptation, provided selection pressures experienced by the different generations are sufficiently similar.

Finally, length of the growing season has ecosystem-level consequences for water, nutrient and carbon cycling. For carbon in particular, it is not obvious whether changes in the length of the growing season will lead to a net increase or decrease in carbon fixation, because of the opposing effects of increases in photosynthesis and respiration. Phenology determines the time period over which photosynthesis can occur, and the increase in primary productivity resulting from this temporal effect can exceed the direct effect of temperature on photosynthetic rate (Piao *et al.* 2007). In this issue, Richardson *et al.* (2010) investigate how this phenological effect on ecosystem productivity varies across temperate forest types and between spring and autumn seasons, showing that an extended growing season can increase net productivity despite increased carbon loss at high temperatures.

Thus, forecasting growing season length under future climate change in various ecosystems is immensely important. However, forecasts of community-level changes in phenology are problematic, given both the rarity of comprehensive long-term datasets and the variability in phenological responses among different species and sites. Ibáñez *et al.* (2010) outline a hierarchical Bayesian approach to this problem that circumvents some of the limitations of more conventional statistical techniques.

As for the consequences of future changes in growing season length, Richardson *et al.* (2010) point out that both spatial proxies and historical conditions are imperfect predictors: changes in species' distributions will interact with phenological changes to affect ecosystem processes (cf. Cleland *et al.* 2007). Indirect effects of growing season length are also possible if, for instance, pest insect outbreaks in longer summers cause severe plant mortality. This suggests a need for incorporating more of the direct and indirect effects of phenology into forecasts of ecosystem change. This entails, in part, knowing the proximate factors regulating phenology and the ultimate factors responsible for current life-history strategies—as discussed earlier. Clearly, this is an enormous challenge, but one that is critical to forecasting the ecological consequences of climate change.

6. EVOLUTION OF PHENOLOGY

Phenology at the population or ecosystem level is ultimately a product of selection acting on variation among individuals. Interest in the evolution of phenological traits such as timing of reproduction or migrations is hardly new, but it has been reinvigorated by recent climate change (e.g. Visser 2008). Even so, our ability to predict how phenologies will evolve in response to recent climate change remains limited. Numerous studies have shown evidence of selection on timing of various biological processes, especially in plants (e.g. Kingsolver *et al.* 2001; Gienapp *et al.* 2006; Elzinga *et al.* 2007; Reed *et al.* 2009), but documented responses to selection in natural environments are rare (Gienapp *et al.* 2008; but see Franks *et al.* 2007). Even in some cases where evolutionary change is expected, based on trait heritabilities and selection pressures, adaptation is not observed (Gienapp *et al.* 2006; van Asch *et al.* 2007), suggesting that we must improve our understanding of the form of selection and constraints on its operation.

In particular, there are some peculiarities to *timing* as a trait that make its evolution especially interesting and challenging to investigate. Adaptive change is facilitated when the trait under selection is also the trait according to which individuals choose mates (Doebeli & Dieckmann 2000; but see Fox 2003). For a trait such as timing of breeding, some level of assortative mating between individuals with similar trait values is inevitable (Fox 2003; Weis & Kossler 2004; Weis 2005), and the resulting inflation of genetic variance can hasten evolutionary change in breeding time (Hendry & Day 2005; Devaux & Lande 2008).

In contrast, the evolutionary lability of phenology can be limited because life-history traits are subject to certain unavoidable constraints. The evidence for genetic correlations limiting the short-term rate of adaptive change is so far surprisingly mixed (Agrawal & Stinchcombe 2009); but basic life-history trade-offs are inescapable, and constraints on the evolution of phenological traits may be relatively widespread (cf. Diggle 1999). In fact, this might explain observations of apparent 'maladaptation' in populations that seem to be frequently mistimed to the phenology of their food sources or mutualists (e.g. Zimmerman *et al.* 1989). As Singer & Parmesan (2010) discuss, such persistent asynchrony may be the result of trade-offs with other, perhaps unmeasured, life-history components (see also Ejsmond *et al.* 2010). Clearly, recognizing the existence of trade-offs and developmental constraints is essential for determining whether current asynchrony in fact represents a negative impact of current climate change—as well as for understanding possible evolutionary responses to future environmental change.

Inherently time-dependent processes such as learning can also influence the evolution of phenology: consumers may take time to learn about the existence or location of a food source, and therefore may ignore individuals of the prey species that appear or reproduce early relative to the population mean. This selective advantage (or disadvantage, if the relationship is mutualistic) to early individuals imposes selection on the relative timing of reproduction, regardless of

the absolute date. At least in theory, this can produce phenological patterns that seem maladaptive at the population level (Forrest & Thomson 2009). In addition, males and females within a population may differ in the optimal timing of emergence or reproduction, because precedence is often favoured in mate competition between males but is less strongly selected in females (Wiklund & Fagerström 1977; Bawa & Beach 1981; Morbey & Ydenberg 2001). This could, in principle, drive sexual conflict over timing (Møller *et al.* 2009), something that could again produce apparently maladaptive features in the population as a whole. Similarly, it has been suggested that protandry could produce negative demographic consequences, and possible 'evolutionary suicide', simply by causing mate-limitation in females when population densities are low (Calabrese & Fagan 2004). These sometimes counterintuitive evolutionary dynamics specific to temporal traits deserve more attention.

There may also be interesting interactions between plasticity and selection on phenological traits. The frequent occurrence of a genetic correlation between the mean value of life-history traits and plasticity in those same traits (Scheiner 1993) complicates the interpretation of selection on phenology (e.g. Nussey *et al.* 2005). In addition, phenotypic plasticity can reduce the strength of selection on the underlying traits, but it can also facilitate adaptation by allowing populations to persist long enough to undergo evolutionary change, or by exposing novel traits on which selection can act (Price *et al.* 2003). Plasticity in the timing of particular life-history stages influences the environmental conditions experienced by, and therefore the nature of selection on, these or later developmental stages (Donohue 2005). So, for example, plastic shifts to earlier flowering or leaf budburst in response to warming temperatures could result in selection for later phenology if early development exposes plants to frost damage. Alternatively, plastic shifts to earlier reproduction in an insect could lead to selection against obligate diapause in offspring if this allowed completion of a second generation per year. Similar ideas about opposing effects of selection and environment on phenotypes have been explored in studies of local adaptation along elevational or latitudinal gradients. For example, high-altitude populations have delayed phenologies relative to their low-elevation counterparts but have been selected for faster development (a phenomenon known as countergradient variation; Conover & Schultz 1995). However, interactions between plastic and genetic changes in phenology have received little study in the context of climate change (though see Crozier *et al.* 2008).

Finally, organisms may have means to 'escape'—either behaviourally or evolutionarily—apparent selection on phenology. For example, seasonal declines in a particular food item might select for increased diet breadth, diet switching or increased dispersal distance instead of earlier phenology. A full, multi-dimensional characterization of the adaptive landscape that would reveal these alternative trajectories will remain an unattainable ideal for most systems. Nevertheless, acknowledging these evolutionary options, as well as the constraints mentioned above, should better allow

us to understand apparent failures to respond to selection on phenology.

7. FUTURE DIRECTIONS

In this introduction, we have tried to give a broad overview of the mechanisms governing phenology and the reasons why phenology is an important factor in evolutionary and ecological research. The articles in this special issue develop many of these topics further, and we hope the issue as a whole will stimulate more synthetic work in this field.

In particular, we perceive some key areas where future research could usefully be focused: first, stronger linkages are needed between the ecology and physiology of phenology. As Visser *et al.* (2010) show, a large body of physiological and chronobiological work relevant to eco-evolutionary studies of phenology has gone largely unnoticed by ecologists and evolutionary biologists because we consult different journals and use different terminologies. Awareness of the linkages between these fields should improve the mechanistic understanding of phenology and forecasts of climate change impacts. The articles by Wilczek *et al.* (2010) and Chuine (2010) illustrate the utility of taking a mechanistic approach to fundamental ecological questions (see also de Senerpont Domis *et al.* (2007) for an application in a different system). Furthermore, deeper knowledge of the developmental and physiological aspects of phenology should improve our understanding of the prospects for evolutionary change in phenological traits (cf. Metcalf & Mitchell-Olds 2009; Singer & Parmesan 2010). This too will be an important component of forecasts of climate change impacts on communities.

Second, there is a need for more information about population-level consequences of phenological variation. An increasing number of documented instances of apparent asynchrony between interacting species leads to obvious questions about the impacts of this asynchrony on the vital rates of the populations involved. Too often, it is impossible to answer these questions because we do not know the baseline degree of synchrony—weather conditions were variable even before recent accelerated climate change, and occasional mismatches must have arisen—and because we do not know how the specific interaction affects population growth rates (the work of Both *et al.* (2006) is a notable exception). It is possible that 'mismatched' species or individuals can often switch to other food sources or move elsewhere. As discussed by Miller-Rushing *et al.* (2010), the population biology of phenology is an area where much work remains to be done.

Third, we argue that an explicit recognition of phenology and seasonality will make for more realistic models of community and ecosystem processes and the ecological impacts of climate change. It is not enough to know the effects of mean annual temperatures and precipitation; we must also know the effects of timing of temperature anomalies and precipitation events. For instance, warming restricted to the cold season may have little impact on populations compared with summer warming, provided winter

temperatures remain below a certain threshold (e.g. Yamanaka *et al.* 2008). Models that incorporate seasonal changes in conditions can reach qualitatively different conclusions than those that assume constant, equilibrium conditions (Steiner *et al.* 2009). Although this may seem an obvious point, most models of community dynamics still operate on the assumption of invariant, or randomly varying, environmental conditions. Similarly, many forecasts of climate-driven changes in species ranges ignore phenology (but see Chuine 2010). Forecasts that incorporate ecological information, particularly niche-based models (e.g. Araújo & Luoto 2007; Wiens *et al.* 2009), tend to focus on whether interacting species will occur in the same place, but neglect to consider whether the temporal aspect of their interactions will be disrupted. Including phenology in these forecasts could yield important insights into future species distributions and interactions.

Because of the interdisciplinary nature of phenology, and the ubiquity of phenological responses to climate change, there are many opportunities for novel synthetic research. Furthermore, timing, as a biological phenomenon, is unique. Unlike other variables, time is not only directional but completely asymmetric: early events can affect later ones, but not vice versa. A plant that germinates and bolts early in a season can change light conditions for its later neighbours, potentially yielding a competitive advantage. As climate conditions and season lengths continue to change, these temporal relationships will also evolve. We expect that the articles collected here will advance our understanding of these changes and point the way for future research.

We thank James Thomson and Elizabeth Wolkovich for their thoughtful comments on the manuscript. J.F. was supported by a scholarship from IODE, Canada.

REFERENCES

- Abrams, P. A. & Rowe, L. 1996 The effects of predation on the age and size of maturity of prey. *Evolution* **50**, 1052–1061. (doi:10.2307/2410646)
- Agrawal, A. F. & Stinchcombe, J. R. 2009 How much do genetic covariances alter the rate of adaptation? *Proc. R. Soc. B* **276**, 1183–1191. (doi:10.1098/rspb.2008.1671)
- Altermatt, F. 2010 Climatic warming increases voltinism in European butterflies and moths. *Proc. R. Soc. B* **277**, 1281–1287. (doi:10.1098/rspb.2009.1910)
- Aono, Y. & Kazui, K. 2008 Phenological data series of cherry tree flowering in Kyoto, Japan, and its application to reconstruction of springtime temperatures since the 9th century. *Int. J. Climatol.* **28**, 905–914. (doi:10.1002/joc.1594)
- Araújo, M. B. & Luoto, M. 2007 The importance of biotic interactions for modelling species distributions under climate change. *Global Ecol. Biogeogr.* **16**, 743–753. (doi:10.1111/j.1466-8238.2007.00359.x)
- Augspurger, C. K. 1981 Reproductive synchrony of a tropical shrub: experimental studies on effects of pollinators and seed predators in *Hybanthus prunifolius* (Violaceae). *Ecology* **62**, 775–788. (doi:10.2307/1937745)
- Ausín, I., Alonso-Blanco, C. & Martínez-Zapater, M. 2005 Environmental regulation of flowering. *Int. J. Dev. Biol.* **49**, 689–705. (doi:10.1387/ijdb.052022ia)
- Balasubramanian, S. *et al.* 2006 The PHYTOCHROME C photoreceptor gene mediates natural variation in flowering and growth responses of *Arabidopsis thaliana*. *Nat. Genet.* **38**, 711–715. (doi:10.1038/ng1818)
- Bauer, S., Gienapp, P. & Madsen, J. 2008 The relevance of environmental conditions for departure decision changes en route in migrating geese. *Ecology* **89**, 1953–1960. (doi:10.1890/07-1101.1)
- Bawa, K. S. & Beach, J. H. 1981 Evolution of sexual systems in flowering plants. *Ann. MO Bot. Gard.* **68**, 254–274. (doi:10.2307/2398798)
- Beatley, J. C. 1974 Phenological events and their environmental triggers in Mojave desert ecosystems. *Ecology* **55**, 856–863. (doi:10.2307/1934421)
- Beebee, T. J. C. 1995 Amphibian breeding and climate. *Nature* **374**, 219–220. (doi:10.1038/374219a0)
- Bosch, J. & Kemp, W. P. 2003 Effect of wintering duration and temperature on survival and emergence time in males of the orchard pollinator *Osmia lignaria* (Hymenoptera: Megachilidae). *Environ. Entomol.* **32**, 711–716. (doi:10.1603/0046-225X-32.4.711)
- Bosch, J. & Kemp, W. P. 2004 Effect of pre-wintering and wintering temperature regimes on weight loss, survival, and emergence time in the mason bee *Osmia cornuta* (Hymenoptera: Megachilidae). *Apidologie* **35**, 469–479. (doi:10.1051/apido:2004035)
- Bostock, J. & Riley, H. T. (eds) 1855 *The natural history of Pliny*. London, UK: H.G. Bohn.
- Both, C., Bouwhuis, S., Lessells, C. M. & Visser, M. E. 2006 Climate change and population declines in a long-distance migratory bird. *Nature* **441**, 81–83. (doi:10.1038/nature04539)
- Both, C., van Asch, M., Bijlsma, R. G., van den Burg, A. B. & Visser, M. E. 2009 Climate change and unequal phenological changes across four trophic levels: constraints or adaptations? *J. Anim. Ecol.* **78**, 73–83. (doi:10.1111/j.1365-2656.2008.01458.x)
- Bradley, N. L., Leopold, A. C., Ross, J. & Huffaker, W. 1999 Phenological changes reflect climate change in Wisconsin. *Proc. Natl Acad. Sci. USA* **96**, 9701–9704. (doi:10.1073/pnas.96.17.9701)
- Bradshaw, W. E. & Holzapfel, C. M. 2001 Genetic shift in photoperiod response correlated with global warming. *Proc. Natl Acad. Sci. USA* **98**, 14 509–14 511. (doi:10.1073/pnas.241391498)
- Bradshaw, W. E. & Holzapfel, C. M. 2006 Evolutionary response to rapid climate change. *Science* **312**, 1477–1478. (doi:10.1126/science.1127000)
- Brearley, F. Q., Proctor, J., Suriantata, Nagy, L., Dalrymple, G. & Voysey, B. C. 2007 Reproductive phenology over a 10-year period in a lowland evergreen rain forest of central Borneo. *J. Ecol.* **95**, 828–839. (doi:10.1111/j.1365-2745.2007.01258.x)
- Brown, R. F. & Mayer, D. G. 1988 Representing cumulative germination. 2. The use of the Weibull function and other empirically derived curves. *Ann. Bot.* **61**, 127–138.
- Buckler, E. S. *et al.* 2009 The genetic architecture of maize flowering time. *Science* **325**, 714–718. (doi:10.1126/science.1174276)
- Calabrese, J. M. & Fagan, W. F. 2004 Lost in time, lonely, and single: reproductive asynchrony and the Allee effect. *Am. Nat.* **164**, 25–37.
- Chuine, I. 2010 Why does phenology drive species distribution? *Phil. Trans. R. Soc. B* **365**, 3149–3160. (doi:10.1098/rstb.2010.0142)
- Clausen, J., Keck, D. D. & Hiesey, W. M. 1941 Regional differentiation in plant species. *Am. Nat.* **75**, 231–250.
- Cleland, E. E., Chiariello, N. R., Loarie, S. R., Mooney, H. A. & Field, C. B. 2006 Diverse responses of phenology to global changes in a grassland ecosystem. *Proc. Natl Acad.*

- Sci. USA* **103**, 13 740–13 744. (doi:10.1073/pnas.0600815103)
- Cleland, E. E., Chuine, I., Menzel, A., Mooney, H. A. & Schwartz, M. D. 2007 Shifting plant phenology in response to global change. *Trends Ecol. Evol.* **22**, 357–365. (doi:10.1016/j.tree.2007.04.003)
- Cloudsley-Thompson, J. L. 1991 *Ecophysiology of desert arthropods and reptiles*. Berlin, Germany: Springer-Verlag.
- Conover, D. O. & Schultz, E. T. 1995 Phenotypic similarity and the evolutionary significance of countergradient variation. *Trends Ecol. Evol.* **10**, 248–252. (doi:10.1016/S0169-5347(00)89081-3)
- Corbet, P. S. 1954 Seasonal regulation in British dragonflies. *Nature* **174**, 655–655. (doi:10.1038/174655a0)
- Crick, H. Q. P. & Sparks, T. H. 1999 Climate change related to egg-laying trends. *Nature* **399**, 423–424. (doi:10.1038/20839)
- Crozier, L. G., Hendry, A. P., Lawson, P. W., Quinn, T. P., Mantua, N. J., Battin, J., Shaw, R. G. & Huey, R. B. 2008 Potential responses to climate change in organisms with complex life histories: evolution and plasticity in Pacific salmon. *Evol. Appl.* **1**, 252–270. (doi:10.1111/j.1752-4571.2008.00033.x)
- Danforth, B. N. 1999 Emergence dynamics and bet hedging in a desert bee, *Perdita portalis*. *Proc. R. Soc. Lond. B* **266**, 1985–1994. (doi:10.1098/rspb.1999.0876)
- Danks, H. V. 2006 Key themes in the study of seasonal adaptations in insects. II. Life-cycle patterns. *Appl. Entomol. Zool.* **41**, 1–13. (doi:10.1303/aez.2006.1)
- Davis, C. C., Willis, C. G., Primack, R. B. & Miller-Rushing, A. J. 2010 The importance of phylogeny to the study of phenological response to global climate change. *Phil. Trans. R. Soc. B* **365**, 3201–3213. (doi:10.1098/rstb.2010.0130)
- de Senerpont Domis, L. N., Mooij, W. M., Hülsmann, S., van Nes, E. H. & Scheffer, M. 2007 Can overwintering versus diapausing strategy in *Daphnia* determine match-mismatch events in zooplankton–algae interactions? *Oecologia* **150**, 682–698. (doi:10.1007/s00442-006-0549-2)
- Devaux, C. & Lande, R. 2008 Incipient allochronic speciation due to non-selective assortative mating by flowering time, mutation and genetic drift. *Proc. R. Soc. B* **275**, 2723–2732. (doi:10.1098/rspb.2008.0882)
- Diekmann, M. 1996 Relationship between flowering phenology of perennial herbs and meteorological data in deciduous forests of Sweden. *Can. J. Bot.* **74**, 528–537.
- Diggle, P. K. 1999 Heteroblasty and the evolution of flowering phenologies. *Int. J. Plant Sci.* **160**(Suppl.), S123–S134.
- Doebeli, M. & Dieckmann, U. 2000 Evolutionary branching and sympatric speciation caused by different types of ecological interactions. *Am. Nat.* **156**, S77–S101.
- Donohue, K. 2005 Niche construction through phenological plasticity: life history dynamics and ecological consequences. *New Phytol.* **166**, 83–92. (doi:10.1111/j.1469-8137.2005.01357.x)
- Dunne, J. A., Harte, J. & Taylor, K. J. 2003 Subalpine meadow flowering phenology responses to climate change: integrating experimental and gradient methods. *Ecol. Monogr.* **73**, 69–86. (doi:10.1890/0012-9615(2003)073[0069:SMFPRT]2.0.CO;2)
- Durant, J. M., Hjermand, D. Ø, Ottersen, G. & Stenseth, N. C. 2007 Climate and the match or mismatch between predator requirements and resource availability. *Climate Res.* **33**, 271–283. (doi:10.3354/cr033271)
- Eilertsen, H. C., Sandberg, S. & Tøllefsen, H. 1995 Photo-periodic control of diatom spore growth: a theory to explain the onset of phytoplankton blooms. *Mar. Ecol. Prog. Ser.* **116**, 303–307. (doi:10.3354/meps116303)
- Ejmsmond, M. J., Czarnoński, M., Kapustka, F. & Kozłowski, J. 2010 How to time growth and reproduction during the vegetative season: an evolutionary choice for indeterminate growers in seasonal environments. *Am. Nat.* **175**, 551–563. (doi:10.1086/651589)
- Ellebjer, S. M., Tamstorf, M. P., Illeris, L., Michelsen, A. & Hansen, B. U. 2008 Inter-annual variability and controls of plant phenology and productivity at Zackenberg. *Adv. Ecol. Res.* **40**, 249–273. (doi:10.1016/S0065-2504(07)00011-6)
- Elzinga, J. A., Atlan, A., Biere, A., Gigord, L., Weis, A. E. & Bernasconi, G. 2007 Time after time: flowering phenology and biotic interactions. *Trends Ecol. Evol.* **22**, 432–439. (doi:10.1016/j.tree.2007.05.006)
- Embree, D. G. 1970 The diurnal and seasonal pattern of hatching of winter moth eggs, *Operophtera brumata* (Geometridae: Lepidoptera). *Can. Entomol.* **102**, 759–768. (doi:10.4039/Ent102759-6)
- Etterson, J. R. & Shaw, R. G. 2001 Constraint to adaptive evolution in response to global warming. *Science* **294**, 151–154. (doi:10.1126/science.1063656)
- Fitter, A. H. & Fitter, R. S. R. 2002 Rapid changes in flowering time in British plants. *Science* **296**, 1689–1691. (doi:10.1126/science.1071617)
- Fitter, A. H., Fitter, R. S. R., Harris, I. T. B. & Williamson, M. H. 1995 Relationships between first flowering date and temperature in the flora of a locality in central England. *Funct. Ecol.* **9**, 55–60. (doi:10.2307/2390090)
- Forrest, J. & Thomson, J. D. 2009 Pollinator experience, neophobia and the evolution of flowering time. *Proc. R. Soc. B* **276**, 935–943. (doi:10.1098/rspb.2008.1434)
- Forrest, J. & Thomson, J. D. 2010 Consequences of variation in flowering time within and among individuals of *Mertensia fusiformis* (Boraginaceae), an early spring wildflower. *Am. J. Bot.* **97**, 38–48. (doi:10.3732/ajb.0900083)
- Forrest, J., Inouye, D. W. & Thomson, J. D. 2010 Flowering phenology in subalpine meadows: does climate variation influence community co-flowering patterns? *Ecology* **91**, 431–440. (doi:10.1890/09-0099.1)
- Fox, G. A. 2003 Assortative mating and plant phenology: evolutionary and practical consequences. *Evol. Ecol. Res.* **5**, 1–18.
- Franks, S. J. & Weis, A. E. 2008 A change in climate causes rapid evolution of multiple life-history traits and their interactions in an annual plant. *J. Evol. Biol.* **21**, 1321–1334. (doi:10.1111/j.1420-9101.2008.01566.x)
- Franks, S. J., Sim, S. & Weis, A. E. 2007 Rapid evolution of flowering time by an annual plant in response to a climate fluctuation. *Proc. Natl Acad. Sci. USA* **104**, 1278–1282. (doi:10.1073/pnas.0608379104)
- Garner, W. W. & Allard, H. A. 1920 Effect of the relative length of day and night and other factors of the environment on growth and reproduction in plants. *J. Agric. Res.* **18**, 553–606.
- Geber, M. A. & Griffen, L. R. 2003 Inheritance and natural selection on functional traits. *Int. J. Plant Sci.* **164**, S21–S42. (doi:10.1086/368233)
- Gienapp, P., Postma, E. & Visser, M. E. 2006 Why breeding time has not responded to selection for earlier breeding in a songbird population. *Evolution* **6**, 2381–2388.
- Gienapp, P., Teplitsky, C., Alho, J. S., Mills, J. A. & Merilä, J. 2008 Climate change and evolution: disentangling environmental and genetic responses. *Mol. Ecol.* **17**, 167–178. (doi:10.1111/j.1365-294X.2007.03413.x)
- Gilbert, J. J. 1974 Dormancy in rotifers. *Trans. Am. Microsc. Soc.* **93**, 490–513. (doi:10.2307/3225154)
- Gillooly, J. F., Charnov, E. L., West, G. B., Savage, V. M. & Brown, J. H. 2002 Effects of size and temperature on developmental time. *Nature* **417**, 70–73. (doi:10.1038/417070a)

- Goldman, B. D. 2001 Mammalian photoperiodic system: formal properties and neuroendocrine mechanisms of photoperiodic time measurement. *J. Biol. Rhythms* **16**, 283–301. (doi:10.1177/074873001129001980)
- Gwinner, E. 1996 Circadian and circannual programmes in avian migration. *J. Exp. Biol.* **199**, 39–48.
- Harrington, R., Woiwod, I. & Sparks, T. 1999 Climate change and trophic interactions. *Trends Ecol. Evol.* **14**, 146–150. (doi:10.1016/S0169-5347(99)01604-3)
- Hegland, S. J., Nielsen, A., Lázaro, A., Bjerknes, A.-L. & Totland, Ø. 2009 How does climate warming affect plant–pollinator interactions? *Ecol. Lett.* **12**, 184–195. (doi:10.1111/j.1461-0248.2008.01269.x)
- Henderson, I. R., Shindo, C. & Dean, C. 2003 The need for winter in the switch to flowering. *Annu. Rev. Genet.* **37**, 371–392. (doi:10.1146/annurev.genet.37.110801.142640)
- Hendry, A. P. & Day, T. 2005 Population structure attributable to reproductive time: isolation by time and adaptation by time. *Mol. Ecol.* **14**, 901–916. (doi:10.1111/j.1365-294X.2005.02480.x)
- Hopkins, A. D. 1918 Periodical events and natural law as guides to agricultural research and practice. *Monthly Weather Rev. Suppl.* **9**, 1–42.
- Høye, T. T. & Forchhammer, M. C. 2008 Phenology of high-arctic arthropods: effects of climate on spatial, seasonal, and inter-annual variation. *Adv. Ecol. Res.* **40**, 299–324. (doi:10.1016/S0065-2504(07)00013-X)
- Hülber, K., Winkler, M. & Grabherr, G. 2010 Intraseasonal climate and habitat-specific variability controls the flowering phenology of high-alpine plant species. *Funct. Ecol.* **24**, 245–252. (doi:10.1111/j.1365-2435.2009.01645.x)
- Husby, A., Kruuk, L. E. B. & Visser, M. E. 2009 Decline in the frequency and benefits of multiple brooding in great tits as a consequence of a changing environment. *Proc. R. Soc. B* **276**, 1845–1854. (doi:10.1098/rspb.2008.1937)
- Ibáñez, I., Primack, R. B., Miller-Rushing, A. J., Ellwood, E., Higuchi, H., Lee, S. D., Kobori, H. & Silander, J. A. 2010 Forecasting phenology under global warming. *Phil. Trans. R. Soc. B* **365**, 3247–3260. (doi:10.1098/rstb.2010.0120)
- Ims, R. A. 1990 The ecology and evolution of reproductive synchrony. *Trends Ecol. Evol.* **5**, 135–140. (doi:10.1016/0169-5347(90)90218-3)
- Inouye, D. W. 2008 Effects of climate change on phenology, frost damage, and floral abundance of montane wildflowers. *Ecology* **89**, 353–362. (doi:10.1890/06-2128.1)
- Jackson, M. T. 1966 Effects of microclimate on spring flowering phenology. *Ecology* **47**, 407–415. (doi:10.2307/1932980)
- Janzen, D. H. 1967 Synchronization of sexual reproduction of trees within the dry season in Central America. *Evolution* **21**, 620–637. (doi:10.2307/2406621)
- Jönsson, A. M., Appelberg, G., Harding, S. & Barring, L. 2009 Spatio-temporal impact of climate change on the activity and voltinism of the spruce bark beetle *Ips typographus*. *Global Change Biol.* **15**, 486–499. (doi:10.1111/j.1365-2486.2008.01742.x)
- Kelly, D. & Sork, V. L. 2002 Mast seeding in perennial plants: why, how, where? *Annu. Rev. Ecol. Syst.* **33**, 427–447. (doi:10.1146/annurev.ecolsys.33.020602.095433)
- Kemp, W. P. & Dennis, B. 1989 Development of two rangeland grasshoppers at constant temperatures: development thresholds revisited. *Can. Entomol.* **121**, 363–371. (doi:10.4039/Ent121363-4)
- Kemp, W. P. & Onsager, J. A. 1986 Rangeland grasshoppers (Orthoptera: Acrididae): modeling phenology of natural populations of six species. *Environ. Entomol.* **15**, 924–930.
- Kemp, W. P., Dennis, B. & Beckwith, R. C. 1986 Stochastic phenology model for the western spruce budworm (Lepidoptera: Tortricidae). *Environ. Entomol.* **15**, 547–554.
- Kimball, S., Angert, A. L., Huxman, T. E. & Venable, D. L. 2010 Contemporary climate change in the Sonoran Desert favors cold-adapted species. *Global Change Biol.* **16**, 1555–1565. (doi:10.1111/j.1365-2486.2009.02106.x)
- Kimberling, D. N. & Miller, J. C. 1988 Effects of temperature on larval eclosion of the winter moth, *Operophtera brumata*. *Entomol. Exp. Appl.* **47**, 249–254. (doi:10.1007/BF00352211)
- Kingsolver, J. G., Hoekstra, H. E., Hoekstra, J. M., Berrigan, D., Vignieri, S. N., Hill, C. E., Hoang, A., Gibert, P. & Beerli, P. 2001 The strength of phenotypic selection in natural populations. *Am. Nat.* **157**, 245–261.
- Laaksonen, T., Ahola, M., Eeva, T., Väisänen, R. A. & Lehikoinen, E. 2006 Climate change, migratory connectivity and changes in laying date and clutch size of the pied flycatcher. *Oikos* **114**, 277–290. (doi:10.1111/j.2006.0030-1299.14652.x)
- Leopold, A. & Jones, S. E. 1947 A phenological record for Sauk and Dane Counties, Wisconsin, 1935–1945. *Ecol. Monogr.* **17**, 81–122. (doi:10.2307/1948614)
- Liedvogel, M., Szulkin, M., Knowles, S. C. L., Wood, M. J. & Sheldon, B. C. 2009 Phenotypic correlates of *Clock* gene variation in a wild blue tit population: evidence for a role in seasonal timing of reproduction. *Mol. Ecol.* **18**, 2444–2456. (doi:10.1111/j.1365-294X.2009.04204.x)
- Logan, J. A., Régnière, J. & Powell, J. A. 2003 Assessing the impacts of global warming on forest pest dynamics. *Front. Ecol. Environ.* **1**, 130–137. (doi:10.1890/1540-9295(2003)001[0130:ATIOWG]2.0.CO;2)
- Masaki, S. 1967 Geographic variation and climatic adaptation in a field cricket (Orthoptera: Gryllidae). *Evolution* **21**, 725–741. (doi:10.2307/2406770)
- Masaki, S. 1980 Summer diapause. *Annu. Rev. Entomol.* **25**, 1–25.
- Mazer, S. J. & LeBuhn, G. 1999 Genetic variation in life-history traits: heritability estimates within and genetic differentiation among populations. In *Life history evolution in plants* (eds T. O. Vuorisalo & P. K. Mutikainen), pp. 85–171. Dordrecht, The Netherlands: Kluwer Academic.
- Medway, L. 1972 Phenology of a tropical rain forest in Malaya. *Biol. J. Linn. Soc.* **4**, 117–146. (doi:10.1111/j.1095-8312.1972.tb00692.x)
- Memmott, J., Craze, P. G., Waser, N. M. & Price, M. V. 2007 Global warming and the disruption of plant–pollinator interactions. *Ecol. Lett.* **10**, 710–717. (doi:10.1111/j.1461-0248.2007.01061.x)
- Metcalf, C. J. E. & Mitchell-Olds, T. 2009 Life history in a model system: opening the black box with *Arabidopsis thaliana*. *Ecol. Lett.* **12**, 593–600. (doi:10.1111/j.1461-0248.2009.01320.x)
- Miller-Rushing, A. J. & Primack, R. B. 2008 Global warming and flowering times in Thoreau's Concord: a community perspective. *Ecology* **89**, 332–341. (doi:10.1890/07-0068.1)
- Miller-Rushing, A. J., Katsuki, T., Primack, R. B., Ishii, Y., Lee, S. D. & Higuchi, H. 2007 Impact of global warming on a group of related species and their hybrids: cherry tree (Rosaceae) flowering at Mt. Takao, Japan. *Am. J. Bot.* **94**, 1470–1478. (doi:10.3732/ajb.94.9.1470)
- Miller-Rushing, A. J., Høye, T. T., Inouye, D. W. & Post, E. 2010 The effects of phenological mismatches on demography. *Phil. Trans. R. Soc. B* **365**, 3177–3186. (doi:10.1098/rstb.2010.0148)
- Mitchell-Olds, T. 1996 Genetic constraints on life-history evolution: quantitative-trait loci influencing growth and

- flowering in *Arabidopsis thaliana*. *Evolution* **50**, 140–145. (doi:10.2307/2410788)
- Møller, A. P., Balbontín, J., Cuervo, J. J., Hermosell, I. G. & de Lope, F. 2009 Individual differences in protandry, sexual selection, and fitness. *Behav. Ecol.* **20**, 433–440. (doi:10.1093/beheco/arn142)
- Morbey, Y. E. & Ydenberg, R. C. 2001 Protandrous arrival timing to breeding areas: a review. *Ecol. Lett.* **4**, 663–673. (doi:10.1046/j.1461-0248.2001.00265.x)
- Morin, X., Lechowicz, M. J., Augspurger, C., O’Keefe, J., Viner, D. & Chuine, I. 2009 Leaf phenology in 22 North American tree species during the 21st century. *Global Change Biol.* **15**, 961–975. (doi:10.1111/j.1365-2486.2008.01735.x)
- Murray, M. B., Cannell, M. G. R. & Smith, R. I. 1989 Date of budburst of fifteen tree species in Britain following climatic warming. *J. Appl. Ecol.* **26**, 693–700.
- Myneni, R. B., Keeling, C. D., Tucker, C. J., Asrar, G. & Nemani, R. R. 1997 Increased plant growth in the northern high latitudes from 1981 to 1991. *Nature* **386**, 698–702. (doi:10.1038/386698a0)
- Nève, G. & Singer, M. C. 2008 Protandry and postandry in two related butterflies: conflicting evidence about sex-specific trade-offs between adult size and emergence time. *Evol. Ecol.* **22**, 701–709.
- Nussey, D. H., Postma, E., Gienapp, P. & Visser, M. E. 2005 Selection on heritable phenotypic plasticity in a wild bird population. *Science* **310**, 304–306. (doi:10.1126/science.1117004)
- Parmesan, C. & Yohe, G. 2003 A globally coherent fingerprint of climate change impacts across natural systems. *Nature* **421**, 37–42. (doi:10.1038/nature01286)
- Paterniani, E. 1969 Selection for reproductive isolation between two populations of maize, *Zea mays* L. *Evolution* **23**, 534–547. (doi:10.2307/2406851)
- Piao, S., Friedlingstein, P., Ciais, P., Viovy, N. & Demarty, J. 2007 Growing season extension and its impact on terrestrial carbon cycle in the Northern Hemisphere over the past 2 decades. *Global Biogeochem. Cycles* **21**, GB3018. (doi:10.1029/2006GB002888)
- Post, E., Pedersen, C., Wilmers, C. C. & Forchhammer, M. C. 2008 Warming, plant phenology and the spatial dimension of trophic mismatch for large herbivores. *Proc. R. Soc. B* **275**, 2005–2013. (doi:10.1098/rspb.2008.0463)
- Price, T., Kirkpatrick, M. & Arnold, S. J. 1988 Directional selection and the evolution of breeding date in birds. *Science* **240**, 798–799. (doi:10.1126/science.3363360)
- Price, T. D., Qvarnström, A. & Irwin, D. E. 2003 The role of phenotypic plasticity in driving genetic evolution. *Proc. R. Soc. Lond. B* **270**, 1433–1440. (doi:10.1098/rspb.2003.2372)
- Rabinowitz, D., Rapp, J. K., Sork, V. L., Rathcke, B. J., Reese, G. A. & Weaver, J. C. 1981 Phenological properties of wind- and insect-pollinated prairie plants. *Ecology* **62**, 49–56. (doi:10.2307/1936667)
- Rathcke, B. & Lacey, E. P. 1985 Phenological patterns of terrestrial plants. *Annu. Rev. Ecol. Syst.* **16**, 179–214. (doi:10.1146/annurev.es.16.110185.001143)
- Réale, D., McAdam, A. G., Boutin, S. & Berteaux, D. 2003 Genetic and plastic responses of a northern mammal to climate change. *Proc. R. Soc. Lond. B* **270**, 591–596. (doi:10.1098/rspb.2002.2224)
- Reed, T. E., Warzybok, P., Wilson, A. J., Bradley, R. W., Wanless, S. & Sydeman, W. J. 2009 Timing is everything: flexible phenology and shifting selection in a colonial seabird. *J. Anim. Ecol.* **78**, 376–387. (doi:10.1111/j.1365-2656.2008.01503.x)
- Régnière, J., Lavigne, D., Dupont, A. & Carter, N. 2007 Predicting the seasonal development of the yellowheaded spruce sawfly (Hymenoptera: Tenthredinidae) in eastern Canada. *Can. Entomol.* **139**, 365–377. (doi:10.4039/N06-054)
- Richardson, A. D. *et al.* 2010 Influence of spring and autumn phenological transitions on forest ecosystem productivity. *Phil. Trans. R. Soc. B* **365**, 3227–3246. (doi:10.1098/rstb.2010.0102)
- Robertson, C. 1924 Phenology of entomophilous flowers. *Ecology* **5**, 393–407. (doi:10.2307/1929302)
- Roff, D. A. 1980 Optimizing development time in a seasonal environment: the ‘ups and downs’ of clinal variation. *Oecologia* **45**, 202–208. (doi:10.1007/BF00346461)
- Rowe, L., Ludwig, D. & Schluter, D. 1994 Time, condition, and the seasonal decline of avian clutch size. *Am. Nat.* **143**, 698–772.
- Roy, D. B. & Sparks, T. H. 2000 Phenology of British butterflies and climate change. *Global Change Biol.* **6**, 407–416. (doi:10.1046/j.1365-2486.2000.00322.x)
- Saino, N., Szép, T., Romano, M., Rubolini, D., Spina, F. & Møller, A. P. 2004 Ecological conditions during winter predict arrival date at the breeding quarters in a trans-Saharan migratory bird. *Ecol. Lett.* **7**, 21–25. (doi:10.1046/j.1461-0248.2003.00553.x)
- Sandrelli, F. *et al.* 2007 A molecular basis for natural selection at the *timeless* locus in *Drosophila melanogaster*. *Science* **316**, 1898–1900. (doi:10.1126/science.1138426)
- Scheiner, S. M. 1993 Genetics and evolution of phenotypic plasticity. *Annu. Rev. Ecol. Syst.* **24**, 35–68. (doi:10.1146/annurev.es.24.110193.000343)
- Schwartz, M. D., Carbone, G. J., Reighard, G. L. & Okie, W. R. 1997 A model to predict peach phenology and maturity using meteorological variables. *Hortscience* **32**, 213–216.
- Sherry, R. A., Zhou, X. H., Gu, S. L., Arnone, J. A., Schimel, D. S., Verburg, P. S., Wallace, L. L. & Luo, Y. Q. 2007 Divergence of reproductive phenology under climate warming. *Proc. Natl Acad. Sci. USA* **104**, 198–202. (doi:10.1073/pnas.0605642104)
- Singer, M. C. & Parmesan, C. 2010 Phenological asynchrony between herbivorous insects and their hosts: signal of climate change or pre-existing adaptive strategy? *Phil. Trans. R. Soc. B* **365**, 3161–3176. (doi:10.1098/rstb.2010.0144)
- Sparks, T. H. & Menzel, A. 2002 Observed changes in seasons: an overview. *Int. J. Climatol.* **22**, 1715–1725. (doi:10.1002/joc.821)
- Sparks, T. H., Jeffree, E. P. & Jeffree, C. E. 2000 An examination of the relationship between flowering times and temperature at the national scale using long-term phenological records from the UK. *Int. J. Biometeorol.* **44**, 82–87. (doi:10.1007/s004840000049)
- Sparks, T. H., Bairlein, F., Bojarinova, J. G., Hüppop, O., Lehtikoinen, E. A., Rainio, K., Sokolov, L. V. & Walker, D. 2005 Examining the total arrival distribution of migratory birds. *Global Change Biol.* **11**, 22–30. (doi:10.1111/j.1365-2486.2004.00887.x)
- Steiner, C. F., Schwaderer, A. S., Huber, V., Klausmeier, C. A. & Litchman, E. 2009 Periodically forced food-chain dynamics: model predictions and experimental validation. *Ecology* **90**, 3099–3107. (doi:10.1890/08-2377.1)
- Steltzer, H., Landry, C., Painter, T. H., Anderson, J. & Ayres, E. 2009 Biological consequences of earlier snowmelt from desert dust deposition in alpine landscapes. *Proc. Natl Acad. Sci. USA* **106**, 11 629–11 634. (doi:10.1073/pnas.0900758106)
- Stinchcombe, J. R., Weinig, C., Ungerer, M., Olsen, K. M., Mays, C., Halldorsdottir, S. S., Purugganan, M. D. & Schmitt, J. 2004 A latitudinal cline in flowering time in *Arabidopsis thaliana* modulated by the flowering time

- gene *FRIGIDA*. *Proc. Natl Acad. Sci. USA* **101**, 4712–4717. (doi:10.1073/pnas.0306401101)
- Stross, R. G. & Hill, J. C. 1968 Photoperiod control of winter diapause in the fresh-water crustacean, *Daphnia*. *Biol. Bull.* **134**, 176–198. (doi:10.2307/1539976)
- Tauber, M. J., Tauber, C. A. & Masaki, S. 1986 *Seasonal adaptations of insects*. New York, NY: Oxford University Press.
- Tauber, E. *et al.* 2007 Natural selection favors a newly derived *timeless* allele in *Drosophila melanogaster*. *Science* **316**, 1895–1898. (doi:10.1126/science.1138412)
- Thomson, J. D. 2010 Flowering phenology, fruiting success and progressive deterioration of pollination in an early-flowering geophyte. *Phil. Trans. R. Soc. B* **365**, 3187–3199. (doi:10.1098/rstb.2010.0115)
- Thórhallsdóttir, T. E. 1998 Flowering phenology in the central highland of Iceland and implications for climatic warming in the Arctic. *Oecologia* **114**, 43–49.
- Tobin, P. C., Nagarkatti, S., Loeb, G. & Saunders, M. C. 2008 Historical projected interactions between climate change insect voltinism in a multivoltine species. *Global Change Biol.* **14**, 951–957. (doi:10.1111/j.1365-2486.2008.01561.x)
- van Asch, M., van Tienderen, P. H., Holleman, L. J. M. & Visser, M. E. 2007 Predicting adaptation of phenology in response to climate change, an insect herbivore example. *Global Change Biol.* **13**, 1596–1604. (doi:10.1111/j.1365-2486.2007.01400.x)
- Van Dijk, H. & Hautekèete, N. 2007 Long day plants and the response to global warming: rapid evolutionary change in day length sensitivity is possible in wild beet. *J. Evol. Biol.* **20**, 349–357. (doi:10.1111/j.1420-9101.2006.01192.x)
- van Noordwijk, A. J. & de Jong, G. 1986 Acquisition and allocation of resources: their influence on variation in life history tactics. *Am. Nat.* **128**, 137–142.
- van Nouhuys, S. & Lei, G. C. 2004 Parasitoid–host meta-population dynamics: the causes and consequences of phenological asynchrony. *J. Anim. Ecol.* **73**, 526–535. (doi:10.1111/j.0021-8790.2004.00827.x)
- van Schaik, C. P., Terborgh, J. W. & Wright, S. J. 1993 The phenology of tropical forests: adaptive significance and consequences for primary consumers. *Annu. Rev. Ecol. Syst.* **24**, 353–377. (doi:10.1146/annurev.es.24.110193.002033)
- Verhulst, S., Tinbergen, J. M. & Daan, S. 1997 Multiple breeding in the great tit. A trade-off between successive reproductive attempts? *Funct. Ecol.* **11**, 714–722. (doi:10.1046/j.1365-2435.1997.00145.x)
- Visser, M. E. 2008 Keeping up with a warming world: assessing the rate of adaptation to climate change. *Proc. R. Soc. B* **275**, 649–659. (doi:10.1098/rspb.2007.0997)
- Visser, M. E., Caro, S. P., van Oers, K., Schaper, S. V. & Helm, B. 2010 Phenology, seasonal timing and circannual rhythms: towards a unified framework. *Phil. Trans. R. Soc. B* **365**, 3113–3127. (doi:10.1098/rstb.2010.0111)
- Wang, R., Farrona, S., Vincent, C., Joecker, A., Schoof, H., Turck, F., Alonso-Blanco, C., Coupland, G. & Albani, M. C. 2009 *PEP1* regulates perennial flowering in *Arabidopsis alpina*. *Nature* **459**, 423–427. (doi:10.1038/nature07988)
- Wayne, N. L. 2001 Regulation of seasonal reproduction in mollusks. *J. Biol. Rhythms* **16**, 391–402. (doi:10.1177/074873001129002097)
- Weis, A. E. 2005 Direct and indirect assortative mating: a multivariate approach to plant flowering schedules. *J. Evol. Biol.* **18**, 536–546. (doi:10.1111/j.1420-9101.2005.00891.x)
- Weis, A. E. & Kossler, T. M. 2004 Genetic variation in flowering time induces phenological assortative mating: quantitative genetic methods applied to *Brassica rapa*. *Am. J. Bot.* **91**, 825–836. (doi:10.3732/ajb.91.6.825)
- Werner, E. E. 1986 Amphibian metamorphosis: growth rate, predation risk, and the optimal size at transformation. *Am. Nat.* **128**, 319–341.
- Wiens, J. A., Stralberg, D., Jongsomjit, D., Howell, C. A. & Snyder, M. A. 2009 Niches, models, and climate change: assessing the assumptions and uncertainties. *Proc. Natl Acad. Sci. USA* **106**, 19 729–19 736. (doi:10.1073/pnas.0901639106)
- Wiklund, C. & Fagerström, T. 1977 Why do males emerge before females? *Oecologia* **31**, 153–158. (doi:10.1007/BF00346917)
- Wilczek, A. M. *et al.* 2009 Effects of genetic perturbation on seasonal life history plasticity. *Science* **323**, 930–934. (doi:10.1126/science.1165826)
- Wilczek, A. M., Burghardt, L. T., Cobb, A. R., Cooper, M. D., Welch, S. M. & Schmitt, J. 2010 Genetic and physiological bases for phenological responses to current and predicted climates. *Phil. Trans. R. Soc. B* **365**, 3129–3147. (doi:10.1098/rstb.2010.0128)
- Williams-Howze, J. 1997 Dormancy in the free-living copepod orders Cyclopoida, Calanoida, and Harpacticoida. *Oceanogr. Mar. Biol. Annu. Rev.* **35**, 257–321.
- Winder, M. & Cloern, J. E. 2010 The annual cycles of phytoplankton biomass. *Phil. Trans. R. Soc. B* **365**, 3215–3226. (doi:10.1098/rstb.2010.0125)
- Wolkovich, E. M. & Cleland, E. E. In press. The phenology of plant invasions: a community ecology perspective. *Front. Ecol. Environ.*
- Yamanaka, T., Tatsuki, S. & Shimada, M. 2008 Adaptation to the new land or effect of global warming? An age-structured model for rapid voltinism change in an alien lepidopteran pest. *J. Anim. Ecol.* **77**, 585–596. (doi:10.1111/j.1365-2656.2008.01367.x)
- Zhang, X., Friedl, M. A. & Schaaf, C. B. 2006 Global vegetation phenology from Moderate Resolution Imaging Spectroradiometer (MODIS): evaluation of global patterns and comparison with *in situ* measurements. *J. Geophys. Res.* **111**, G04017. (doi:10.1029/2006JG000217)
- Zhang, X., Tarpley, D. & Sullivan, J. T. 2007 Diverse responses of vegetation phenology to a warming climate. *Geophys. Res. Lett.* **34**, L19405. (doi:10.1029/2007GL031447)
- Zimmerman, J. K., Roubik, D. W. & Ackerman, J. D. 1989 Asynchronous phenologies of a neotropical orchid and its euglossine bee pollinator. *Ecology* **70**, 1192–1195. (doi:10.2307/1941389)